

ABSTRACT

Any algorithm that applies to all MDPs will suffer $\Omega(\sqrt{|\mathcal{S}||\mathcal{A}|T})$ regret on some MDP. So what do we do when $|\mathcal{S}|, |\mathcal{A}|$ are **extremely large or infinite**? The curse of dimensionality means our only hope is to exploit some low-dimensional structure.

We show that if the MDP can be parameterized within some known function class, we obtain regret **bounds that scale with the dimensionality, rather than cardinality, of the system**. We characterize this dependence explicitly in terms of the eluder dimension. We also present a simple and computationally efficient algorithm (PSRL) that satisfies these bounds. These are the **first regret bounds for general model-based learning**.

PROBLEM FORMULATION

Learn to optimize a random finite horizon MDP M in repeated finite episodes of interaction.



Figure 1: classic reinforcement learning setting

- State space \mathcal{S} , action space \mathcal{A}
- Rewards $r_t \sim R^M(s_t, a_t) \in \mathcal{R}$
- Transitions $s_{t+1} \sim P^M(s_t, a_t) \in \mathcal{P}$
- Episode length τ , define $t_k := (k-1)\tau + 1$

For MDP M and policy μ , define a value function

$$V_{\mu,i}^M(s) := \mathbb{E}_{M,\mu} \left[\sum_{j=i}^{\tau} \bar{R}^M(s_j, a_j) \mid s_i = s \right],$$

Define the regret in episode k using μ_k on M^*

$$\Delta_k := \int_{\mathcal{S}} \rho(s) \left(\underbrace{V_{\mu_k^*,1}^{M^*}(s)}_{\text{optimal value}} - \underbrace{V_{\mu_k,1}^{M^*}(s)}_{\text{actual value}} \right)$$

And finally $\text{Regret}(T, \pi, M^*) := \sum_{k=1}^{\lceil T/\tau \rceil} \Delta_k$.

Naive exploration such as Boltzman or ϵ -greedy can lead to exponential regret. Good performance requires balancing **exploration vs exploitation**.

ELUDER DIMENSION



Eluder principle: a measurement at x is independent of $\{x_1, \dots, x_n\}$ if functions that are similar at $\{x_1, \dots, x_n\}$ could differ significantly at x .

Definition 1 ((\mathcal{F}, ϵ) -dependence).

We will say that $x \in \mathcal{X}$ is (\mathcal{F}, ϵ) -dependent on $\{x_1, \dots, x_n\} \subseteq \mathcal{X} \iff \forall f, \tilde{f} \in \mathcal{F} \subseteq \{f: \mathcal{X} \rightarrow \mathbb{R}^n\}$
 $\sum_{i=1}^n \|f(x_i) - \tilde{f}(x_i)\|_2^2 \leq \epsilon^2 \implies \|f(x) - \tilde{f}(x)\|_2 \leq \epsilon$.
 $x \in \mathcal{X}$ is (ϵ, \mathcal{F}) -independent of $\{x_1, \dots, x_n\}$ iff it does not satisfy the definition for dependence.




Figure 2: x_5 is $(\{f_1, f_2\}, 1)$ -independent of $\{x_1, \dots, x_4\}$.

Definition 2 (Eluder Dimension = $\dim_E(\mathcal{F}, \epsilon)$).

The length of the longest possible sequence of elements in \mathcal{X} such that for some $\epsilon' \geq \epsilon$ every element is (\mathcal{F}, ϵ') -independent of its predecessors.


Examples

- \mathcal{X} finite $\implies \dim_E(\mathcal{F}, \epsilon) \leq |\mathcal{X}|$.
- $\mathcal{F} \subseteq \{f: \mathbb{R}^n \rightarrow \mathbb{R}^p \text{ linear}\}$
 $\implies \dim_E(\mathcal{F}, \epsilon) = O(np \log(1/\epsilon))$

KOLMOGOROV DIMENSION

The Kolmogorov dimension of a function class \mathcal{F} :

$$\dim_K(\mathcal{F}) := \limsup_{\alpha \downarrow 0} \frac{\log(N(\mathcal{F}, \alpha, \|\cdot\|_2))}{\log(1/\alpha)}$$



In this diagram $N(\mathcal{F}, \alpha, \|\cdot\|_2) \leq 7$

Example: $\dim_K(\mathbb{R}^d) = d$

LIPSCHITZ SMOOTHNESS

Definition 3 (Future value function U_i^M).

For any distribution Φ over \mathcal{S} we define:

$$U_i^M(\Phi) := \mathbb{E}_{M,\mu^M} [V_{\mu^M,i+1}^M(s) \mid s \sim \Phi]$$

as the value of the optimal policy, starting from Φ .

- Learning an infinite MDP requires regularity
- Assume $U_i^{M^*}$ is Lipschitz in $\mathbb{E}[s \mid s \sim \Phi]$ wrt $\|\cdot\|_2$
- Satisfied whenever $V_{\mu^*,i}^{M^*}$ Lipschitz in s wrt $\|\cdot\|_2$
- But this is a strictly weaker condition since system noise can help smooth future value.

POSTERIOR SAMPLING

For each episode k :

1. Sample an MDP from the posterior distribution for the true MDP: $M_k \sim \phi(\cdot \mid H_t)$.
2. Use policy $\mu_k \in \arg \max_{\mu} V_{\mu}^{M_k}$.

MAIN RESULTS

If M^* is an MDP with rewards $R^* \in \mathcal{R}$ and transitions $P^* \in \mathcal{P}$ with sub σ -Gaussian noise then the **expected regret** to time T of PSRL is bounded:

$$\tilde{O} \left(\underbrace{\sigma_{\mathcal{R}} \sqrt{d_K(\mathcal{R}) d_E(\mathcal{R}) T}}_{\text{rewards}} + \underbrace{\mathbb{E}[K^*]}_{\text{Lipschitz}} \underbrace{\sigma_{\mathcal{P}} \sqrt{d_K(\mathcal{P}) d_E(\mathcal{P}) T}}_{\text{transitions}} \right)$$

Notation:

- Kolmogorov dimension $d_K(\mathcal{F}) := \dim_K(\mathcal{F})$
- Eluder dimension $d_E(\mathcal{F}) := \dim_E(\mathcal{F}, T^{-1})$
- Lipschitz constant K^* for future value function

Corollary:

Let M^* be a linear-quadratic system in \mathbb{R}^d with σ -sub-Gaussian noise mean-bounded by C then:

$$\mathbb{E}[\text{Regret}(T, \pi^{PS}, M^*)] = \underbrace{\tilde{O}(\sigma C d^2 \sqrt{T})}_{\text{no exponential scaling in } d}$$

REFERENCES

Please see the full paper:
<http://arxiv.org/abs/1406.1853>



PROOF SKETCH

We consider the regret in an episode k :

$$\begin{aligned} \Delta_k &= V_{*,1}^*(s) - V_{k,1}^*(s) \\ &= \underbrace{(V_{k,1}^k(s) - V_{k,1}^*(s))}_{\text{Imagined - Actual}} + \underbrace{(V_{*,1}^*(s) - V_{k,1}^k(s))}_{\mathbb{E}[\cdot]=0 \text{ by posterior}} \end{aligned}$$

We can decompose this into Bellman error:

$$V_{k,1}^k - V_{k,1}^* = \sum_{i=1}^{\tau} \underbrace{(\mathcal{T}_{k,i}^k - \mathcal{T}_{k,i}^*)}_{B := \text{Bellman error}} V_{k,i+1}^k + \sum_{i=1}^{\tau} \underbrace{d_{t_k+1}}_{\mathbb{E}=0 \text{ martingale}}$$

We can now use the Hölder inequality to bound:

$$B \leq \sum_{i=1}^{\tau} \left\{ \underbrace{|\bar{R}^k - \bar{R}^*|}_{\text{reward error}} + \underbrace{K^k}_{\text{Lipschitz}} \underbrace{\|P^k - P^*\|_2}_{\text{transition error}} \right\}$$

We conclude the proof by upper bounding these deviations in terms of our estimation errors on R^* and P^* . We use concentration inequalities to express the error bounds for \mathcal{R} and \mathcal{P} in terms of the eluder dimension and Kolmogorov dimension.

Note a proof for a similar optimistic algorithm is possible, however this would require a generally intractable planning step. We believe that the sampling approach will also be more statistically efficient since it is not affected by loose analysis.

SO WHAT?

- **Practical reinforcement learning problems often have $|\mathcal{S}|$ and $|\mathcal{A}|$ very large or infinite.**
- **“Tabula rasa” learning will always require minimum $T = \Omega(|\mathcal{S}||\mathcal{A}|)$ for good guarantees \implies must exploit low-dimensional structure.**
- **We produce a unified analysis for model-based RL in terms of the dimensionality, rather than the cardinality, of the system.**
- **Conceptually simple, computationally efficient algorithm PSRL satisfies these bounds.**

CONTACT INFORMATION

Web www.stanford.edu/~iosband
Email iosband@stanford.edu

